Available online at www.sciencedirect.com
SCIENCE DIRECT®
ELSEVIER
Discrete Applied Mathematics 149 (2005) 219
DISCRETE APPLIED MATHEMATICS
www.elsevier.com/locate/dam

Author index to volume 149

Bioch, J.C., The complexity of modular decomposition of Boolean functions	(1–3)	1–13
Čepek, O. and P. Kučera, Known and new classes of generalized Horn formulae with polynomial recognition and SAT testing	(1–3)	14–52
Cohen, D., M. Cooper, P. Jeavons and A. Krokhin, Supermodular functions and the complexity of MAX CSP	(1–3)	53–72
Cooper, M., see D. Cohen	(1–3)	53–72
Cusick, T.W. and Y. Li, k-th order symmetric SAC boolean functions and bisecting binomial coefficients	(1–3)	73–86
Faye, A. and Q. Trinh, A polyhedral approach for a constrained quadratic 0–1 problem	(1–3)	87–100
Geist, D., see B. Schieber	(1–3)	154–173
Golumbic, M.C., see A. Mintz	(1–3)	131–153
Hatami, H. and H. Maserrat, On the computational complexity of defining sets	(1–3)	101–110
Jansen, T. and I. Wegener, Real royal road functions—where crossover provably is essential	(1–3)	111–125
Jeavons, P., see D. Cohen	(1–3)	53–72
Krokhin, A., see D. Cohen	(1–3)	53–72
Kučera, P., On the size of maximum renamable Horn sub-CNF	(1–3)	126–130
Kučera, P., see O. Čepek	(1–3)	14–52
Li, Y., see T.W. Cusick	(1–3)	73–86
Maserrat, H., see H. Hatami	(1–3)	101–110
Mintz, A. and M.C. Golumbic, Factoring Boolean functions using graph partitioning	(1–3)	131–153
Schieber, B., D. Geist and A. Zaks, Computing the minimum DNF representation of Boolean functions defined by intervals	(1–3)	154–173
Trinh, Q. see A. Faye	(1–3)	87–100
Wegener, I., see T. Jansen	(1–3)	111–125
Wild, M., The many benefits of putting stack filters into disjunctive or conjunctive normal form	(1–3)	174–191
Xiao, W., Linear symmetries of Boolean functions	(1–3)	192–199
Zaks, A., see B. Schieber	(1–3)	154–173
Zverovich, I.E., Characterizations of closed classes of Boolean functions in terms of forbidden subfunctions and Post classes	(1–3)	200–218

doi:10.1016/S0166-218X(05)00152-6